



Davis Spine & Orthopaedics

Spine and Orthopaedic Surgeon
Board Certified and Fellowship Trained
www.DavisSpine.com
331 Oak Manor Drive Suite 201
Glen Burnie, MD 2106
Phone: 410-625-5050 Fax: 410-766-1404

Scoliosis Corrective Surgery Discharge instructions

When to call 911:

- **If you are having shortness of breath or chest pain, call 911**
- **If you are having swallowing problems that interfere with your ability to breathe call 911 and go to the ER. You MUST also contact the office to make us aware of your trip to the ER, we have no other way to know you are there.**

When to call the office: (even if it is after hours – call before going to the ER)

- **If your temperature exceeds 101.5° F**
- **If your incision begins to separate**
- **If your incision begins to drain more that it did at your time of discharge. If you have drainage that is soaking through your dressing after one hour**
- **If you have an increase in pain, swelling or redness around the incision**
- **If you have difficulty with controlling your bowel or bladder (not constipation, the opposite)**
- **If you have loss of movement in one or both legs**

Discomfort:

- Pain - After surgery, pain is managed with narcotic medication. Because narcotic pain pills are addictive, they are used for a limited period (8 to 12 weeks). Thereafter, pain is managed with acetaminophen (e.g., Tylenol).
- Constipation- Use of narcotics may cause constipation, drink lots of water and eat high fiber foods. Laxatives (e.g., Dulcolax, Senokot, Milk of Magnesia) or glycerin suppositories can also be helpful, all of which can be bought without a prescription.
- Hoarseness, sore throat, or difficulty swallowing may occur in some patients and **should not be cause for alarm**. These symptoms usually resolve in 1 to 4 weeks.
- It is usual to experience some continued back and leg pain. This should progressively improve. Often, cramping pain or soreness is associated with increasing your activities too quickly. If this happens to you, try restricting your activities for a day or so which will often help dramatically. If there is a return of severe pain call the office.

Restrictions:

- **DO NOT** take NSAIDs or steroids (i.e.- prednisone, motrin, ibuprofen, aleve)
- **DO NOT SMOKE!** – This may prevent your spine from fusing and surgery will have been a waste of time, yours and mine!
- **Wear your hard clamshell brace AT ALL TIMES WHILE AWAKE for at least 12 weeks.** You may remove it for sleeping and showers. It will cover your back, chest and hips and should be **tight**.
- Once you progress out of the hard brace, you will wear a soft corset for an additional 12 weeks while awake. It may be removed for showers and sleep.

- Log roll in and out of bed
- Motion restrictions – No trunk flexion, extension, side bending or rotation
- Do not lift, stoop, stretch, reach overhead or bend during the early period following surgery. If you must lift an object, bend at the knees.
- Do not lift anything heavier than 8 pounds (e.g., gallon of milk).
- **Do not** drive for 12 weeks after surgery. You may ride in a car, but avoid long trips until after you have returned for a follow up visit. Never drive while using narcotics.
- Avoid sitting for long periods of time, avoid overstuffed or extremely soft chair. A firm, hardback chair is best, use chairs with arms, it will be easier to get up.
- Housework and yard-work **are not permitted** until you are told otherwise. This includes gardening, mowing, vacuuming, ironing, and loading/unloading the dishwasher, washer, or dryer.
- Postpone sexual activity until your 8 week follow-up appointment. If x-rays look good at this visit, you may resume sexual activity as a passive participant, and while wearing your brace.

Activity:

- Nap if you are tired but **DO NOT** stay in bed all day.
- Use the incentive spirometer (breathing device) provided by the hospital – once each hour while you are awake for 2 weeks after surgery.
- You may need help with daily activities (e.g., dressing, bathing), however, most patients are able to care for themselves right away.
- Gradually return to your normal activities. Walking is encouraged; start with a short distance and gradually increase up to 1 to 2 miles daily.

Bathing/Incision Care:

- You should change the dressing daily for about two weeks. Do **not** apply any creams or ointments for the first 4-6 weeks.
- Sutures are placed under the skin and will be absorbed by your body. There will be small pieces of adhesive tape superficially, which can be removed in about 14 days. If these become dirty or sticky, they may be removed earlier.
- You may shower 5 to 7 days after surgery. The incision may get wet, pat it dry with sterile gauze. Once dry, cover with a dry sterile dressing.
- No tub baths, hot tubs, or swimming pools until you are told it's safe to do so.

Recovery and prevention:

We will see you for follow up 2 weeks after your surgery and again at 8 weeks. Recovery time generally lasts 8 to 12 weeks with improvement each week. We will discuss the type of work you do at your follow up visits and we will decide together when to release you back to work. We will have you bring x-rays to your 8 week follow up to check your progress, then again at your 12 week follow up.

Recurrences of back pain are common. The key to avoiding recurrence is prevention:

- Use proper lifting techniques
- Use good posture during sitting, standing, moving, and sleeping.
- Maintain a healthy weight and lean body mass
- Please---**NO SMOKING!!!**